

AMENDMENTS TO THE SPECIFICATION

Page 8, please replace the first full paragraph with the following amended paragraph:

The term “high glucose medium” as used herein refers to an animal cell culture medium further containing D-glucose ranging from 3,500 to 5,500 ~~mg/l~~ ~~mg/ml~~ and sodium pyruvate ranging from 50 to 200 ~~mg/l~~ ~~mg/ml~~. Representative commercially available high glucose media include, but are not limited to, HG(high glucose)-DMEM, IMDM and so on.

Page 9, please replace the last paragraph that bridges to page 10 with the following amended paragraph:

In order to isolate and culture multipotent progenitor/stem cells from the cord blood-derived mononuclear cells, the mononuclear cells are cultured in a series of animal cell culture media consisting of a first, second and third media in order. The animal cell culture medium used in the present invention is a conventional animal cell culture medium employed in the art which may further contain additional ingredients and/or antibiotics according to the particular purpose of cultivation. Representative commercially available animal cell culture media include, but are not limited to, RPMI1640, MEM, α -MEM, IMDM or DMEM, preferably DMEM. Preferably, the animal cell culture medium is a high glucose medium which further contains D-glucose ranging from 3,500 to 5,500 ~~mg/l~~ ~~mg/ml~~ and sodium pyruvate ranging from 50 to 200 ~~mg/l~~ ~~mg/ml~~ and may contain additional ingredients and/or antibiotics according to the particular purpose of cultivation. In a preferred embodiment of the present invention, HG-DMEM (Gibco

Cat. No. 12800-017) containing additional 4,500 ~~mg/l~~ ~~mg/ml~~ of D-glucose and 110 ~~mg/l~~ ~~mg/ml~~ of sodium pyruvate is employed as the high glucose medium.

Page 10, please replace the last paragraph that bridges to page 11 with the following amended paragraph:

After the second culture, when mono-layer cell colony forming cells are grown to 80 to 90% of adhesion, the culture medium is removed, and the cells are washed with PBS and treated with trypsin/EDTA, to recover the cells. The cells are inoculated into the third animal cell culture medium at a concentration ranging from 2×10^4 to 8×10^4 cells/cm² and cultured at 37°C under an atmosphere of 5% CO₂ for 1 to 2 weeks to induce growth and proliferation of the cells into multipotent progenitor/stem cells. Preferably, the culture medium is replaced with a fresh medium at an interval of 3 to 5 days. The cultivation in the third animal cell culture medium is to maintain undifferentiated state of the cells cultured in the second animal cell culture medium and induce proliferation thereof. At this time, the third animal cell culture medium is the same as the first animal cell culture medium except that G-CSF is replaced with SCF and EGF. Preferably, the third animal cell culture medium is a high glucose culture medium further containing 10 to 20% FBS, 1 to 2 mM L-glutamine, 10 to 100 ~~ng/ml~~ ~~ng/ml~~ of SCF and 5 to 50 ~~ng/ml~~ ~~ng/ml~~ of EGF. SCF and EGF induce the proliferation of stem cells, and in particular, SCF plays the role of maintaining the multipotent progenitor/stem cells.